



Review

Contributions of estrogen receptor- α and estrogen receptor- β to the regulation of behaviorMarc J. Tetel^a, Donald W. Pfaff^{b,*}^a Neuroscience Program, Wellesley College, Wellesley, MA, USA^b Laboratory of Neurobiology and Behavior, The Rockefeller University, New York, NY, USA

ARTICLE INFO

Article history:

Received 28 May 2009

Received in revised form 15 January 2010

Accepted 16 January 2010

Available online 25 January 2010

Keywords:

Estrogen receptor

Hypothalamus

Preoptic area

Lordosis

Aggression

Progesterin receptor

Sexual differentiation

Steroid receptor coactivator-1 (SRC-1)

ABSTRACT

Studies of the mechanisms by which estrogens influence brain function and behavior have advanced from the explication of individual hormone receptors, neural circuitry and individual gene expression. Now, we can report patterns of estrogen receptor subtype contributions to patterns of behavior. Moreover, new work demonstrates important contributions of nuclear receptor coactivator expression in the central nervous system. In this paper, our current state of knowledge is reviewed.

© 2010 Published by Elsevier B.V.

Gene products coding for nuclear hormone receptors have afforded neuroscientists unusual opportunities for the analysis of brain mechanisms that regulate behavior. Brain researchers benefit from the relatively simple chemistry of steroid hormones and from the large numbers of reagents effective in the manipulation of hormone-dependent systems of cells. Success during the recent scientific generations of work in this field has allowed us to understand how sex steroid actions in the brain coordinate behavioral regulation with other factors in the body and in the environment. The upshot is that we understand mechanisms with physical dimensions ranging from Angstrom units (ligand binding domains of estrogen receptors) to light years (the seasonality of reproduction in many animals).

Of particular interest are gene duplication products among steroid hormone receptors, both of which the ancestral hormone can activate both as the normal ligand, but which are genetically and chemically distinguishable. Estrogens elicit many of their effects on behavior and physiology by binding to its intracellular receptors, estrogen receptor- α (ER α) and ER β , which are transcribed from different genes [1,2] and are members of the steroid/nuclear receptor superfamily of transcriptional activators [3,4]. While estrogens can function in the brain by binding to receptors on the cell membrane and rapidly activate cytoplasmic signaling pathways [5–8], this review will focus on the

effects of estrogens on behavior and physiology mediated by ER α and ER β via classic, genomic mechanisms of action.

Using mice created by homologous recombination to achieve null mutations for ER α (ER α KO) or ER β (ER β KO), we and others have conducted a large number of behavioral and histochemical assays in genetic females and genetic males. Several of the assays simply depend on the classical ER, without a discernible contribution from ER β . During other assays, expression of ER β appears to act as a brake on ER α function. In still other assays, ER β and ER α can substitute for each other.

The contributions of ER α and ER β to behaviors and physiological events are complicated and in no way uniform. Below we have summarized results gathered to date, and labeled each according to the relative contributions of the ER subtypes. The extreme scenarios are that ER α and ER β act in an obligatory synergistic fashion $\{\alpha + \beta\}$ or that they exert equal and opposite function $\{\alpha$ vs. $\beta\}$. The data suggest that different patterns of neural function require different patterns of contributions by ER α and ER β , such that: ER α and ER β can substitute for each other $\{\alpha$ or $\beta\}$; the contributions of ER α and ER β are not related $\{\alpha/\beta$ contributions nr}; ER β acts as a brake on ER α function and reduces the effects of ER α $\{\alpha - \beta = \alpha'\}$. While we and others have used many behavioral or histochemical assays, the references below concentrate on data related directly to the brain's regulation of reproduction. Then, our main inference is stated at the end of this paper. For an in depth review of ER α and ER β expression in brain regions known to regulate behaviors associated with

* Corresponding author.

E-mail address: pfaff@mail.rockefeller.edu (D.W. Pfaff).

reproduction, including sexual behavior, aggression and anxiety, please see Bodo et al. [9].

1. Behavior

Perhaps the most striking result was the first reported: that putting an ER α KO female into the cage of a sexually experienced male had two surprising results: (i) that she behaves exactly as a male would, with respect to immediate aggressive behavior, and (ii) that she is treated like an intruder male by the resident male [10]. In a broader framework, comparisons among results of the different labs cited below attest to the reliability of neuroendocrine and behavioral results in this field.

1.1. Female sexual behavior $\{\alpha - \beta = \alpha'\}$

Gene expression from ER α raises levels of generalized arousal in female mice [11] eventuating in significantly higher locomotor activity [12], important for the initiation of courtship behaviors. In turn, studies using ER α KO mice and RNA interference indicate that the ER α gene product is essential for the normal performance of lordosis behavior [13–16]. In contrast, while ER β KO females are sub-fertile [17], their lordosis behavior was equal to that of wt females and somewhat elevated in that they exhibited strong sexual receptivity the day after estrus [14,16]. These results indicate that female sexual behavior is dependent on the classical ER α , and not ER β which appears to reduce the effects of ER α on sexual receptivity.

1.2. Male sexual behavior $\{\alpha/\beta$ contributions nr}

While adult ER β KO males exhibit normal male sex behavior in regard to mounts, intromissions and ejaculations [16,18], this ER subtype appears to be important for development of male sex behavior. Peripubertal ER β KO males display their first ejaculations at a later age than their wt littermates [19]. In adults, reduced success in progress toward ejaculation is seen in ER α KO males [20]; that is, latencies through the series of intromissions are increased so that the temporal interval from first intromission to eventual ejaculation is increased – ejaculation is delayed. Finally, knocking out both ER α and ER β completely abolishes male sexual behaviors, including mounting. [21]. Thus it appears that both ER subtypes contribute to the normal display of male sexual behavior. We infer from these data that even under circumstances in which ER β by itself is not necessary for male sex behavior, it still can add synergistically to ER α , such that the double knockout completely abolishes males sexual behaviors.

1.3. Aggression in male ERKO mice $\{\alpha - \beta = \alpha'\}$

Early work has shown that ER α KO males are less aggressive to a male intruder than wt males [20,22]. Another study found that while wt males treated with testosterone (T) only attack T-treated male intruders, ER α KO males display equal aggression towards T-treated male, and estradiol (E)-treated female, intruders [23]. This inability of ER α KO males to distinguish intruders, possibly due to impaired processing of chemosensory information, suggests that ER α is important in normal social preferences. Furthermore, gonadally intact ER β KO males exhibit higher levels of aggression compared to wt males [20]. In gonadectomized males, estradiol benzoate (EB) treatment induces higher levels of aggression in ER β KO animals than wt [24]. These data suggest that the effects of estrogens on male aggression are stimulated by ER α and suppressed by ER β .

1.4. Anxiety in females $\{\alpha - \beta = \alpha'\}$

A variety of studies indicate that ovarian hormones influence anxiety in female rodents [25–27]. To test the function of the two ER subtypes in

anxiety, ovariectomized rats were given DPN (diarylpropionitrile, an ER β selective agonist), PPT (Tris(4-hydroxyphenyl)-4-propyl-1H-pyrazole, an ER α selective agonist), E or vehicle and tested for anxiety-related behaviors [28]. DPN treatment decreased anxiety-like behaviors in elevated plus maze and open field tests. Anxiogenic behaviors, including the time spent grooming, were decreased by DPN and increased by PPT [28]. It should be pointed out that the selectivity of DPN and PPT for ER β and ER α , respectively, are not absolute [29–31]. However, in support of this anxiolytic effect of ER β , ER β KO females display higher levels of anxiety than wt littermates by spending less time exploring the distal portion of the open arm of a maze [32,33]. This effect in ovariectomized ER β KO mice was observed whether mice were treated with chronic E or vehicle [32]. Taken together, these findings indicate that ER β is important during development and/or adulthood in mediating the anxiolytic effects of estrogens on anxiety.

1.5. Stress in male rats $\{\alpha/\beta$ contributions nr}

The paraventricular nucleus (PVN) of the hypothalamus is critical in the modulation of stress and expresses high levels of androgen receptors (AR) [34] and ER β [35–37], but very little ER α [35]. To examine the role of ER β and AR in the regulation of the hypothalamic–pituitary–adrenal response to stress, gonadectomized male rats were implanted with pellets containing E, dihydrotestosterone (DHT, which binds AR and is not aromatized to estradiol), 5 α -androstane-3 β ,17 β -diol (3 β -diol, which binds ER β with a higher affinity than ER α ; [29], DPN or PPT into the PVN [38]. Following restraint stress, animals treated with DHT, 3 β -diol or DPN had reduced levels of stress-induced release of corticosterone and ACTH, while E or PPT treatment increased the release of these stress hormones. These data suggest that ER β functions to decrease HPA activity, while stimulation of ER α enhances it. The ER antagonist, tamoxifen, partially blocked the effects of 3 β -diol and DHT, while an AR antagonist did not. The authors suggest that these effects are mediated through ER β , since 3 β -diol binds ER β with a higher affinity than ER α [29]. In addition, administration into the PVN of DHT, 3 β -diol or DPN reduced the restraint-induced increase in *c-fos* mRNA in this brain region [38]. These findings suggest that ER β inhibits the HPA axis in males and that DHT may suppress the stress response, in part, via conversion to 3 β -diol and its subsequent binding to ER β .

2. Sexual differentiation of the brain

The gonadal steroids elicit profound organizational effects on the developing brain that result in adult sexually dimorphic brain physiology and behavior. During the perinatal period, the testes secrete testosterone that masculinizes (increased adult male sexual behavior) and defeminizes (decreased adult female sexual behavior) adult reproductive behavior. In the brain, testosterone can be converted to DHT or estradiol, which binds to AR and ER, respectively. Thus, testosterone can ultimately have both androgenic and estrogenic effects on the developing and adult brain.

2.1. Sexual differentiation of dopamine neurons in the anteroventral periventricular area $\{\alpha$ or β

In the hypothalamus of rodents, the anteroventral periventricular area (AVPV) has a sexually dimorphic subpopulation of dopaminergic neurons that is larger in females than males and is thought to be involved in the normal display of the preovulatory surge [39]. Testicular feminized male (tfm) mice, which have disrupted AR, and wt males had equal numbers of tyrosine-hydroxylase (TH)-immunoreactive neurons (indicative of dopamine neurons) in the AVPV, suggesting that AR do not regulate the expression of dopamine neurons in this brain region [40]. However, ER α KO male mice had more TH-immunoreactive neurons than wt males, indicating that

disruption of the ER α gene feminized the number of dopaminergic neurons in the AVPV [40]. In an extension of this work, both ER α KO and ER β KO male mice had levels of TH-immunoreactive neurons that were higher than wt males and equal to wt female mice [41]. In addition, treatment of wt female mice during postnatal days 1–3 with either estradiol, DPN or PPT reduced the number of TH-immunoreactive neurons in the AVPV [41]. Taken together, these studies suggest that both ER α and ER β , and not AR, are important in the sexual differentiation of the dopaminergic system in the AVPV of male and female mice.

3. Estrogen-responsive genes

3.1. Estradiol-induction of PR: females $\{\alpha|\beta$ contributions nr}, males $\{\alpha + \beta\}$, and developing female brain $\{\alpha - \beta = \alpha'\}$

A classic example of an estrogen-responsive gene is the estradiol-induction of PR in a variety of tissues, including brain. While PR are present in low levels in the brains of ovariectomized rodents, E-priming dramatically increases the expression of PR in the medial preoptic area (MPOA), ventromedial hypothalamus (VMH), arcuate and midbrain central gray [42–52]. Interestingly, the developing rat brain is extremely sensitive to estradiol, given that doses as low as 100 ng of E are sufficient to induce PR [53]. In adult female guinea pigs virtually all of the E-induced PR cells express ER α [49,54], while about 30% of the E-induced PR cells in rat brain express ER β [37].

In females, it appears that ER α is the major contributor to E-induction of PR in the brain. In support, E-induction of PR in the VMH and POA of ER β KO females was equal to wt animals [14,55], suggesting an important role for ER α in these regions. However, while E-induction of PR is greatly diminished in ER α KO female mice, it is not abolished in all areas, including the POA and VMH [14,56,57]. This residual induction of PR by estradiol may be due to ER β (in the absence of ER α) or ER α splice variants that have been detected in ER α KO mice [56]. In contrast to females, while E treatment did not induce PR expression in the VMH and POA of wt males, it did increase PR levels in both brain regions of ER β KO males [55]. These findings suggest an important role for ER β in the sexually dimorphic responses to estrogens in reproductively-relevant brain regions. In addition, these data suggest that ER β can modulate the responsiveness of ER α in some cells.

Emilie Rissman's lab has used heterozygous breeders to generate mice with varying numbers of functional and disrupted ER genes to determine the contribution of each ER subtype to E-induction of PR in the MPOA [58]. Male and female mice with either 0, 1 or 2 functional copies of ER α and/or ER β were gonadectomized and received implants of estradiol or blank capsules for 5 days. In females, maximal induction of PR by E in the MPOA required at least one functional copy of the ER α gene. In contrast to females, one copy of each ER α and ER β genes are required for the full induction of PR in the male MPOA. These data support the concept that the role of ER β in PR induction is sexually dimorphic and that there is functional dependence of the two ERs in PR induction in the male brain.

These findings in the adult brain suggest that ER β can alter the activity of ER α . This concept is supported by a study in developing rat brain using ER subtype selective agonists [59]. In the developing female ventromedial nucleus of the hypothalamus (VMN), selective activation of ER α with PPT induces PR to higher levels than estradiol alone, which activates both ER subtypes. In contrast, in the MPN, which expresses high levels of ER α and low amounts of ER β [60], PPT and estradiol induced PR to the same extent. In addition, selective activation of ER β by DPN did not induce PR in the MPN. Moreover, activation of ER β diminishes the ER α -mediated increase of PR in the VMN, but not the MPN. These findings suggest that activation of ER β by estradiol in the VMN inhibits ER α transcriptional activity, and thus suppresses ER α -mediated transactivation of the PR gene in this brain

region [59]. This suppressive effect of ER β may provide a mechanism to protect the VMN from defeminization by estradiol during important periods of development, thus ensuring proper feminization of the VMN and subsequent adult female sexual behavior. This idea is supported by work in mice showing that activation of ER α or ER β during development alters female sexual behavior in adulthood [61].

3.2. Estrogen and androgen regulation of AVP $\{\alpha|\beta$ contributions nr}

Arginine vasopressin (AVP), a neurotransmitter involved in aggression, is regulated by androgens and estrogens. Testosterone, and to a lesser extent estradiol, up-regulates AVP expression in brain regions involved in aggression [62,63]. To investigate the roles of ER α and AR in gonadal hormone regulation of AVP and aggression, mice lacking a functional AR (tfm) were compared with ER α KO mice crossed with tfm (resulting in a double AR/ER α KO), ER α KO and wt mice [64]. Gonadectomized mice were treated with estradiol and analyzed by resident-intruder aggression tests. Wt and tfm males displayed aggressive behaviors, while ER α KO males and AR/ER α KO males did not, suggesting that ER α functions in males to regulate aggression as discussed above. AVP levels in the lateral septum revealed that wt and tfm males had the highest levels, followed by ER α KO males, and lowest levels in wt females and AR/ER α KO males. AR/ER α KO males expressed the lowest levels of AVP in the medial amygdala, with all other genotypes expressing equal amounts in this region [64]. In contrast to the lateral septum, estradiol reduces AVP expression in the male PVN [65]. This down-regulation of AVP by estradiol is abolished in ER β KO males, indicating that ER β is critical for normal AVP expression in this brain region [65]. Interestingly, another study using a neuronal cell line found that DHT, and its metabolite androstane-3 β ,17 β -diol, bind to ER β to stimulate AVP promoter activity [66]. It will be essential for future studies to more fully investigate the function of ER β in AVP expression in the brain. Taken together, these findings indicate that both ER α and AR are required for maximal expression of AVP in the lateral septum and medial amygdala, while ER β is critical in AVP expression in the PVN.

4. ER α and ER β interactions and mechanisms

In order to better understand the functional differences of ER α and ER β , one must understand the mechanisms of action of these two subtypes. The ER subtypes have high homology in the DNA binding domain (97%), but much lower homology in the AF-1 and ligand binding domain (60%) [67]. These structural differences allow for variation in the binding affinity and specificity for different ligands of the two ER subtypes [29,68]. In addition, these differences in ligand binding, and the subsequent distinct changes in receptor conformation, enable the ER subtypes to vary in their ability to interact with different estrogen response elements [69]. Thus, ER α and ER β can have profoundly different effects in the transduction of estrogen signaling.

4.1. ER α and ER β interactions with nuclear receptor coactivators

In addition to binding to different ligands, ER α and ER β may elicit distinct transcriptional effects through differential interactions with nuclear receptor coactivators. Nuclear receptor coactivators enhance steroid receptor transcriptional activity *in vitro* [70,71]. Moreover, these coactivators have been found to be critical in estrogen action in brain and behavior [72,73]. For example, steroid receptor coactivator-1 (SRC-1), a member of the p160 family of coactivators, is critical for maximal ER-mediated transactivation of the PR gene in the VMN [74,75] and the display of estrogen-dependent behaviors [76]. Recent studies using pull-down assays have explored the physical interactions between the ER subtypes and SRC-1 from female rat brains [77]. SRC-1 from hypothalamus or hippocampus interacted with ER α and ER β when bound to estradiol, which was confirmed by mass

spectrometry. SRC-1 may function with ER α in the hypothalamus to mediate expression of female sexual behavior [13–16], and with both ER subtypes in the hippocampus to differentially modulate estrogen's effects on cognition [9,78] and stress [9,79].

The different functions of the ER subtypes in the brain may be explained in part by the lower transcriptional activity of ER β observed in particular cell lines [80]. These differences in transcriptional abilities between ER α and ER β may be attributed to differential recruitment of coactivators, or differences in the ability of the same coactivator to facilitate transcription of the ER subtypes [81]. Interestingly, while SRC-1 from the hippocampus interacted equally with the ER subtypes, SRC-1 from hypothalamus interacted more with ER α than with ER β . These findings suggest that ER α is a more efficient transcriptional activator of SRC-1 dependent signaling pathways in the hypothalamus than ER β [77]. In support, previous work indicates that SRC-1 function in the hypothalamus is important for normal expression of ER-mediated female sexual receptivity [76], which as discussed above is ER α -dependent [13–16,82]. These differential interactions of SRC-1 from hypothalamus or hippocampus with the ER subtypes suggest that these brain regions have distinct expression patterns of coregulators involved in these important protein–protein interactions. In addition, it is possible that SRC-1 undergoes differential phosphorylation in these two brain regions, leading to distinct patterns of interaction with the ER subtypes. These findings suggest the importance of using biologically-relevant tissue, in contrast to the use of cell lines alone, in investigating receptor–coactivator interactions. It may be that other cofactors and proteins that are present in tissue, such as the brain, are important for appropriate coactivator interactions with receptor. Understanding how nuclear receptor coactivators function with various steroid receptors, and their subtypes, is critical to understanding how hormones act in different brain regions to profoundly influence physiology and behavior.

It has been suggested that the lack of ER β results in ER β KO mice that are hypersensitive to E [55]. In support, E induces elevated levels of PR in the brains of ER β KO males [55] and the uteri of ER β KO females [83]. In addition, sexual behavior and learning studies suggest that ER β KO females are hypersensitive to E [16,78]. ER β repression of ER α activity may involve the sequestration of shared nuclear receptor coactivators, that is relieved due to the absence of ER β in ER β KO mice.

4.2. DNA as an allosteric ligand

Another level of ER signaling regulation may be exerted by DNA on the ER subtypes. In a recent elegant series of studies, Meijsing et al. [84] demonstrate that DNA can act as an allosteric ligand of glucocorticoid receptors (GR). Different GR binding sequences, which are imperfect palindromic, hexameric half sites separated by 3-base pair spacers, that differ by as little as one base pair, can elicit distinct conformations in GR and regulate its transcriptional activity. As suggested by the authors, this idea of DNA sequences acting as allosteric ligands may modulate signaling of other nuclear receptors, including ER [84]. In support, the AF-2 domain of ER binds to different coregulatory peptides when ER is bound to distinct DNA sequences [69,85]. Thus, in addition to the type of ligand that is available to ER, specific DNA sequences that bind the ER subtypes may modulate the conformation of ER α and/or ER β in the cell and influence the coregulators that it recruits, thus altering the transcriptional activity of the receptors.

5. Conclusions

Taken together, the complex dependencies of results in this field of neuroendocrinology serve to defeat simplistic thinking about gene/behavior relationships. Even with these clear genetic manipulations and straightforward behavioral assays, unexpected results have

emerged. For example, with respect to the impact of an ER β KO on aggressive behaviors by male mice, it became clear that the strength of the phenotype depends upon the age at which the male is tested, the KO effect being strongest just after puberty [86]. Even more surprising, the relation between a given gene and a given behavior (in this case, aggressive behavior) can be exactly the opposite between males and females (summarized in [87]). In addition, given the importance of coactivators in steroid receptor signaling, it is important to consider the possible effects on the signaling of one receptor subtype when the other subtype is knocked out. For example, consider a population of cells that coexpress ER α and ER β [37]. In ER α KO mice, it is possible that in this population of cells, ER β signaling may be enhanced due to decreased competition for common coactivators now that ER α has been eliminated. While we are not aware of any published data supporting this idea, it would be important to consider and explore these potential cell-specific effects in future studies.

We note that the prize-winning discovery of George Beadle and Edward Tatum, based on their work in 1941 with the fungus *Neurospora*, depended on their enunciation of the “one gene – one enzyme” concept. As distinct from their conclusion, and as inferred in Pfaff et al., 2002 [88], we believe that the strategic advantages of studying steroid hormone action in the brain have allowed us to state, as modern neuroscientists, that *patterns* of gene expression govern *patterns* of behavioral functions.

To ultimately understand the differential role of these ER subtypes on complex behaviors, it will be critical to identify and explore the different molecules that function in ER transcription in the brain. Elucidating the myriad of modifications of chromatin elicited by these coactivators and other regulatory molecules in brain will be essential. Moreover, understanding how these proteins assemble at the promoter, in a brain region and cell-type specific fashion, will be an exciting challenge for neuroendocrinologists to tackle in the near future.

Acknowledgements

Studies contributed by the authors' laboratories were supported by grants from National Science Foundation IBN 0080818 and National Institutes of Health R01 DK61935 (MJT) and NIH MH-38273 and HD-05751 (DWP).

References

- [1] E.V. Jensen, T. Suzuki, T. Kawasima, W.E. Stumpf, P.W. Jungblut, E.R. de Sombre, A two-step mechanism for the interaction of estradiol with rat uterus, *Proc. Natl. Acad. Sci. U. S. A.* 59 (1968) 632–638.
- [2] G.G.J.M. Kuiper, E. Enmark, M. Peltö-Huikko, S. Nilsson, J. Gustafsson, Cloning of a novel estrogen receptor expressed in rat prostate and ovary, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 5925–5930.
- [3] D.J. Mangelsdorf, C. Thummel, M. Beato, P. Herrlich, G. Schütz, K. Umesono, B. Blumberg, P. Kastner, M. Mark, P. Chambon, R.M. Evans, The nuclear receptor superfamily: the second decade, *Cell* 83 (1995) 835–839.
- [4] M.J. Tsai, B.W. O'Malley, Molecular mechanisms of action of steroid/thyroid receptor superfamily members, *Ann. Rev. Biochem.* 63 (1994) 451–486.
- [5] N. Vasudevan, D.W. Pfaff, Non-genomic actions of estrogens and their interaction with genomic actions in the brain, *Front. Neuroendocrinol.* 29 (2008) 238–257.
- [6] P. Micevych, J. Kuo, A. Christensen, Physiology of membrane oestrogen receptor signalling in reproduction, *J. Neuroendocrinol.* 21 (2009) 249–256.
- [7] P.G. Mermelstein, Membrane-localised oestrogen receptor alpha and beta influence neuronal activity through activation of metabotropic glutamate receptors, *J. Neuroendocrinol.* 21 (2009) 257–262.
- [8] M.J. Kelly, O.K. Ronnekleiv, Membrane-initiated estrogen signaling in hypothalamic neurons, *Mol. Cell. Endocrinol.* 290 (2008) 14–23.
- [9] C. Bodo, E.F. Rissman, New roles for estrogen receptor beta in behavior and neuroendocrinology, *Front. Neuroendocrinol.* 27 (2006) 217–232.
- [10] S. Ogawa, J.A. Taylor, D.B. Lubahn, K.S. Korach, D.W. Pfaff, Reversal of sex roles in genetic female mice by disruption of estrogen receptor gene, *Neuroendocrinology* 64 (1996) 467–470.
- [11] J. Garey, A. Goodwillie, J. Frohlich, M. Morgan, J.A. Gustafsson, O. Smithies, K.S. Korach, S. Ogawa, D.W. Pfaff, Genetic contributions to generalized arousal of brain and behavior, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 11019–11022.

- [12] S. Ogawa, J. Chan, J.A. Gustafsson, K.S. Korach, D.W. Pfaff, Estrogen increases locomotor activity in mice through estrogen receptor alpha: specificity for the type of activity, *Endocrinology* 144 (2003) 230–239.
- [13] S. Ogawa, V. Eng, J. Taylor, D.B. Lubahn, K.S. Korach, D.W. Pfaff, Roles of estrogen receptor-alpha gene expression in reproduction-related behaviors in female mice, *Endocrinology* 139 (1998) 5070–5081.
- [14] A.E. Kudwa, E.F. Rissman, Double oestrogen receptor alpha and beta knockout mice reveal differences in neural oestrogen-mediated progesterin receptor induction and female sexual behaviour, *J. Neuroendocrinol.* 15 (2003) 978–983.
- [15] S. Musatov, W. Chen, D.W. Pfaff, M.G. Kaplitt, S. Ogawa, RNAi-mediated silencing of estrogen receptor alpha in the ventromedial nucleus of hypothalamus abolishes female sexual behaviors, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 10456–10460.
- [16] S. Ogawa, J. Chan, A.E. Chester, J.A. Gustafsson, K.S. Korach, D.W. Pfaff, Survival of reproductive behaviors in estrogen receptor beta gene- deficient (betaERKO) male and female mice, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 12887–12892.
- [17] J.H. Kregel, J.B. Hodgins, J.F. Couse, E. Enmark, M. Warner, J.F. Mahler, M. Sar, K.S. Korach, J. Gustafsson, O. Smithies, Generation and reproductive phenotypes of mice lacking estrogen receptor-beta, *Proc. Natl. Acad. Sci. U.S.A.* 95 (1998) 15677–15682.
- [18] A.E. Kudwa, C. Bodo, J.A. Gustafsson, E.F. Rissman, A previously uncharacterized role for estrogen receptor beta: defeminization of male brain and behavior, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 4608–4612.
- [19] J.L. Temple, E.M. Scordalakes, C. Bodo, J.A. Gustafsson, E.F. Rissman, Lack of functional estrogen receptor beta gene disrupts pubertal male sexual behavior, *Horm. Behav.* 44 (2003) 427–434.
- [20] S. Ogawa, D.B. Lubahn, K.S. Korach, D.W. Pfaff, Behavioral effects of estrogen receptor gene disruption in male mice, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 1476–1481.
- [21] S. Ogawa, A.E. Chester, S.C. Hewitt, V.R. Walker, J.A. Gustafsson, O. Smithies, K.S. Korach, D.W. Pfaff, Abolition of male sexual behaviors in mice lacking estrogen receptors alpha and beta (alpha beta ERKO), *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 14737–14741.
- [22] S. Ogawa, T.F. Washburn, J. Taylor, D.B. Lubahn, K.S. Korach, D.W. Pfaff, Modifications of testosterone-dependent behaviors by estrogen receptor-alpha gene disruption in male mice, *Endocrinology* 139 (1998) 5058–5069.
- [23] E.M. Scordalakes, E.F. Rissman, Aggression in male mice lacking functional estrogen receptor alpha, *Behav. Neurosci.* 117 (2003) 38–45.
- [24] M. Nomura, S. Andersson, K.S. Korach, J.A. Gustafsson, D.W. Pfaff, S. Ogawa, Estrogen receptor-beta gene disruption potentiates estrogen-inducible aggression but not sexual behaviour in male mice, *Eur. J. Neurosci.* 23 (2006) 1860–1868.
- [25] M.A. Morgan, D.W. Pfaff, Estrogen's effects on activity, anxiety, and fear in two mouse strains, *Behav. Brain Res.* 132 (2002) 85–93.
- [26] K.M. Frick, L.A. Burlingame, J.A. Arters, J. Berger-Sweeney, Reference memory, anxiety and estrous cyclicity in C57BL/6NIA mice are affected by age and sex, *Neuroscience* 95 (2000) 293–307.
- [27] A.Y. Galeeva, S.G. Pivina, P. Tuohimaa, N.E. Ordyan, Involvement of nuclear progesterone receptors in the formation of anxiety in female mice, *Neurosci. Behav. Physiol.* 37 (2007) 843–848.
- [28] T.D. Lund, T. Rovis, W.C. Chung, R.J. Handa, Novel actions of estrogen receptor-beta on anxiety-related behaviors, *Endocrinology* 146 (2005) 797–807.
- [29] G.G. Kuiper, J.G. Lemmen, B. Carlsson, J.C. Corton, S.H. Safe, P.T. van der Saag, D.B. van, J.A. Gustafsson, Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta, *Endocrinology* 139 (1998) 4252–4263.
- [30] S.R. Stauffer, C.J. Coletta, R. Tedesco, G. Nishiguchi, K. Carlson, J. Sun, B.S. Katzenellenbogen, J.A. Katzenellenbogen, Pyrazole ligands: structure-affinity/activity relationships and estrogen receptor-alpha-selective agonists, *J. Med. Chem.* 43 (2000) 4934–4947.
- [31] W.R. Harrington, S. Sheng, D.H. Barnett, L.N. Petz, J.A. Katzenellenbogen, B.S. Katzenellenbogen, Activities of estrogen receptor alpha- and beta-selective ligands at diverse estrogen responsive gene sites mediating transactivation or transrepression, *Mol. Cell. Endocrinol.* 206 (2003) 13–22.
- [32] D.B. Imwalle, J.A. Gustafsson, E.F. Rissman, Lack of functional estrogen receptor beta influences anxiety behavior and serotonin content in female mice, *Physiol. Behav.* 84 (2005) 157–163.
- [33] A.A. Walf, C.J. Koonce, C.A. Frye, Estradiol or diarylpropionitrile decrease anxiety-like behavior of wildtype, but not estrogen receptor beta knockout, mice, *Behav. Neurosci.* 122 (2008) 974–981.
- [34] L. Zhou, J.D. Blaustein, G.J. De Vries, Distribution of androgen receptor immunoreactivity in vasopressin- and oxytocin-immunoreactive neurons in the male rat brain, *Endocrinology* 134 (1994) 2622–2627.
- [35] P.J. Shughrue, M.V. Lane, I. Merchenthaler, Comparative distribution of estrogen receptor-alpha and -beta mRNA in the rat central nervous system, *J. Comp. Neurol.* 388 (1997) 507–525.
- [36] S.E. Alves, V. Lopez, B.S. McEwen, N.G. Weiland, Differential colocalization of estrogen receptor beta (ERbeta) with oxytocin and vasopressin in the paraventricular and supraoptic nuclei of the female rat brain: an immunocytochemical study, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 3281–3286.
- [37] B. Greco, E.A. Allegretto, M.J. Tetel, J.D. Blaustein, Coexpression of ER beta with ER alpha and progesterin receptor proteins in the female rat forebrain: effects of estradiol treatment, *Endocrinology* 142 (2001) 5172–5181.
- [38] T.D. Lund, L.R. Hinds, R.J. Handa, The androgen 5alpha-dihydrotestosterone and its metabolite 5alpha-androstan-3beta, 17beta-diol inhibit the hypothalamo-pituitary-adrenal response to stress by acting through estrogen receptor beta-expressing neurons in the hypothalamus, *J. Neurosci.* 26 (2006) 1448–1456.
- [39] R.B. Simerly, L.W. Swanson, R.A. Gorski, The distribution of monoaminergic cells and fibers in a periventricular preoptic nucleus involved in the control of gonadotropin release: immunohistochemical evidence for a dopaminergic sexual dimorphism, *Brain Res.* 330 (1985) 55–64.
- [40] R.B. Simerly, M.C. Zee, J.W. Pendleton, D.B. Lubahn, K.S. Korach, Estrogen receptor-dependent sexual differentiation of dopaminergic neurons in the preoptic region of the mouse, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 14077–14082.
- [41] C. Bodo, A.E. Kudwa, E.F. Rissman, Both estrogen receptor-alpha and -beta are required for sexual differentiation of the anteroventral periventricular area in mice, *Endocrinology* 147 (2006) 415–420.
- [42] N.J. MacLusky, B.S. McEwen, Oestrogen modulates progesterin receptor concentrations in some rat brain regions but not in others, *Nature* 274 (1978) 276–278.
- [43] M. Warembourg, Radioautographic study of the rat brain, uterus and vagina after [³H]R-5020 injection, *Mol. Cell Endocrinol.* 12 (1978) 67–79.
- [44] N.J. MacLusky, B.S. McEwen, Progesterin receptors in rat brain: distribution and properties of cytoplasmic progesterin-binding sites, *Endocrinology* 106 (1980) 192–202.
- [45] J.D. Blaustein, H.H. Feder, Cytoplasmic progesterin receptors in female guinea pig brain and their relationship to refractoriness in expression of female sexual behavior, *Brain Res.* 177 (1979) 489–498.
- [46] A.H. Lauber, G.J. Romano, D.W. Pfaff, Sex difference in estradiol regulation of progesterin receptor messenger RNA in rat mediobasal hypothalamus as demonstrated by *In situ* hybridization, *Neuroendocrinology* 53 (1991) 608–613.
- [47] S. Ogawa, U.E. Olazabal, I.S. Parhar, D.W. Pfaff, Effects of intrahypothalamic administration of antisense DNA for progesterone receptor mRNA on reproductive behavior and progesterone receptor immunoreactivity in female rat, *J. Neurosci.* 14 (1994) 1766–1774.
- [48] J.D. Blaustein, J.C. King, D.O. Toft, J. Turcotte, Immunocytochemical localization of estrogen-induced progesterin receptors in guinea pig brain, *Brain Res.* 474 (1988) 1–15.
- [49] M. Warembourg, A. Jolivet, E. Milgrom, Immunohistochemical evidence of the presence of estrogen and progesterone receptors in the same neurons of the guinea pig hypothalamus and preoptic area, *Brain Res.* 480 (1989) 1–15.
- [50] C. Beyer, N. Damm, V. Brito, E. Kupperts, Developmental expression of progesterone receptor isoforms in the mouse midbrain, *Neuroreport* 13 (2002) 877–880.
- [51] R.E.M. Scott, X.S. Wu-Peng, D.W. Pfaff, Regulation and expression of progesterone receptor mRNA isoforms A and B in the male and female rat hypothalamus and pituitary following oestrogen treatment, *J. Neuroendocrinol.* 14 (2002) 175–183.
- [52] R.D. Brinton, R.F. Thompson, M.R. Foy, M. Baudry, J. Wang, C.E. Finch, T.E. Morgan, C.J. Pike, W.J. Mack, F.Z. Stanczyk, J. Nilsen, Progesterone receptors: form and function in brain, *Front. Neuroendocrinol.* 29 (2008) 313–339.
- [53] W.C. Chung, T.R. Pak, M.J. Weiser, L.R. Hinds, M.E. Andersen, R.J. Handa, Progesterin receptor expression in the developing rat brain depends upon activation of estrogen receptor alpha and not estrogen receptor beta, *Brain Res.* 1082 (2006) 50–60.
- [54] J.D. Blaustein, J.C. Turcotte, Estradiol-induced progesterin receptor immunoreactivity is found only in estrogen receptor-immunoreactive cells in guinea pig brain, *Neuroendocrinology* 49 (1989) 454–461.
- [55] J.L. Temple, H.N. Fugger, X. Li, S.J. Shetty, J. Gustafsson, E.F. Rissman, Estrogen receptor beta regulates sexually dimorphic neural responses to estradiol, *Endocrinology* 142 (2001) 510–513.
- [56] C.A. Moffatt, E.F. Rissman, M.A. Shupnik, J.D. Blaustein, Induction of progesterin receptors by estradiol in the forebrain of estrogen receptor-alpha gene-disrupted mice, *J. Neurosci.* 18 (1998) 9556–9563.
- [57] P.J. Shughrue, D.B. Lubahn, A. Negro-Vilar, K.S. Korach, I. Merchenthaler, Responses in the brain of estrogen receptor alpha-disrupted mice, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 11008–11012.
- [58] A.E. Kudwa, J.A. Gustafsson, E.F. Rissman, Estrogen receptor B modulates estradiol induction of progesterin receptor immunoreactivity in male, but not in female, mouse medial preoptic area, *Endocrinology* 145 (2004) 4500–4506.
- [59] K.L. Gonzales, M.J. Tetel, C.K. Wagner, Estrogen receptor (ER) beta modulates ERalpha responses to estrogens in the developing rat ventromedial nucleus of the hypothalamus, *Endocrinology* 149 (2008) 4615–4621.
- [60] S.E. Perez, E.Y. Chen, E.J. Mufson, Distribution of estrogen receptor alpha and beta immunoreactive profiles in the postnatal rat brain, *Brain Res. Dev. Brain Res.* 145 (2003) 117–139.
- [61] A.E. Kudwa, V. Michopoulos, J.D. Gatewood, E.F. Rissman, Roles of estrogen receptors alpha and beta in differentiation of mouse sexual behavior, *Neuroscience* 138 (2006) 921–928.
- [62] G.J. De Vries, B.J. Crenshaw, H.A. Al-Shamma, Gonadal steroid modulation of vasopressin pathways, *Ann. N. Y. Acad. Sci.* 652 (1992) 387–396.
- [63] G.J. De Vries, Z. Wang, N.A. Bullock, S. Numan, Sex differences in the effects of testosterone and its metabolites on vasopressin messenger RNA levels in the bed nucleus of the stria terminalis of rats, *J. Neurosci.* 14 (1994) 1789–1794.
- [64] E.M. Scordalakes, E.F. Rissman, Aggression and arginine vasopressin immunoreactivity regulation by androgen receptor and estrogen receptor alpha, *Genes Brain Behav.* 3 (2004) 20–26.
- [65] M. Nomura, E. McKenna, K.S. Korach, D.W. Pfaff, S. Ogawa, Estrogen receptor-beta regulates transcript levels for oxytocin and arginine vasopressin in the hypothalamic paraventricular nucleus of male mice, *Brain Res. Mol. Brain Res.* 109 (2002) 84–94.
- [66] T.R. Pak, W.C. Chung, L.R. Hinds, R.J. Handa, Estrogen receptor-(beta) mediates dihydrotestosterone-induced stimulation of the arginine vasopressin promoter in neuronal cells, *Endocrinology* 148 (2007) 3371–3382.
- [67] G.G. Kuiper, E. Enmark, M. Peltö-Huikko, S. Nilsson, J.A. Gustafsson, Cloning of a novel receptor expressed in rat prostate and ovary, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 5925–5930.

- [68] G.G.J.M. Kuiper, B. Carlsson, K. Grandien, E. Enmark, J. Häggblad, S. Nilsson, J. Gustafsson, Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta, *Endocrinology* 138 (1997) 863–870.
- [69] J.M. Hall, D.P. McDonnell, K.S. Korach, Allosteric regulation of estrogen receptor structure, function, and coactivator recruitment by different estrogen response elements, *Mol. Endocrinol.* 16 (2002) 469–486.
- [70] M.G. Rosenfeld, V.V. Lunyak, C.K. Glass, Sensors and signals: a coactivator/corepressor/epigenetic code for integrating signal-dependent programs of transcriptional response, *Genes Dev.* 20 (2006) 1405–1428.
- [71] B.W. O'Malley, Molecular biology. Little molecules with big goals, *Science* 313 (2006) 1749–1750.
- [72] M.J. Tetel, Nuclear receptor coactivators: essential players in steroid hormone action in brain and behavior, *J. Neuroendocrinol.* 21 (2009) 229–237.
- [73] M.J. Tetel, A.P. Auger, T.D. Charlier, Who's in charge? Nuclear receptor coactivator and corepressor function in brain and behavior, *Front. Neuroendocrinol.* 30 (2009) 328–342.
- [74] H.A. Molenda, A.L. Griffin, A.P. Auger, M.M. McCarthy, M.J. Tetel, Nuclear receptor coactivators modulate hormone-dependent gene expression in brain and female reproductive behavior in rats, *Endocrinology* 143 (2002) 436–444.
- [75] E.M. Apostolakis, M. Ramamurphy, D. Zhou, S. Onate, B. O'Malley, Acute disruption of select steroid receptor coactivators prevents reproductive behavior in rats and unmasks genetic adaptation in knockout mice, *Mol. Endocrinol.* 16 (2002) 1511–1523.
- [76] H.A. Molenda-Figueira, C.A. Williams, A.L. Griffin, E.M. Rutledge, J.D. Blaustein, M.J. Tetel, Nuclear receptor coactivators function in estrogen receptor- and progesterin receptor-dependent aspects of sexual behavior in female rats, *Horm. Behav.* 50 (2006) 383–392.
- [77] H.A. Molenda-Figueira, S.D. Murphy, K.L. Shea, N.K. Siegal, Y. Zhao, J.G. Chadwick, L.A. Denner, M.J. Tetel, Steroid receptor coactivator-1 from brain physically interacts differentially with steroid receptor subtypes, *Endocrinology* 149 (2008) 5272–5279.
- [78] H.N. Fugger, T.C. Foster, J. Gustafsson, E.F. Rissman, Novel effects of estradiol and estrogen receptor alpha and beta on cognitive function, *Brain Res.* 883 (2000) 258–264.
- [79] C. Isgor, M. Cecchi, M. Kabbaj, H. Akil, S.J. Watson, Estrogen receptor beta in the paraventricular nucleus of hypothalamus regulates the neuroendocrine response to stress and is regulated by corticosterone, *Neuroscience* 121 (2003) 837–845.
- [80] F. Delaunay, K. Pettersson, M. Tujague, J.A. Gustafsson, Functional differences between the amino-terminal domains of estrogen receptors alpha and beta, *Mol. Pharmacol.* 58 (2000) 584–590.
- [81] C. Wong, B. Komm, B.J. Cheskis, Structure–function evaluation of ER alpha and beta interplay with SRC family coactivators. ER selective ligands, *Biochemistry* 40 (2001) 6756–6765.
- [82] E.F. Rissman, A.H. Early, J.A. Taylor, K.S. Korach, D.B. Lubahn, Estrogen receptors are essential for female sexual receptivity, *Endocrinology* 138 (1997) 507–510.
- [83] Z. Weihua, S. Saji, S. Makinen, G. Cheng, E.V. Jensen, M. Warner, J.A. Gustafsson, Estrogen receptor (ER) beta, a modulator of ERalpha in the uterus, *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 5936–5941.
- [84] S.H. Meijjing, M.A. Pufall, A.Y. So, D.L. Bates, L. Chen, K.R. Yamamoto, DNA binding site sequence directs glucocorticoid receptor structure and activity, *Science* 324 (2009) 407–410.
- [85] D.M. Heery, E. Kalkhoven, S. Hoare, M.G. Parker, A signature motif in transcriptional co-activators mediates binding to nuclear receptors, *Nature* 387 (1997) 733–736.
- [86] M. Nomura, L. Durbak, J. Chan, O. Smithies, J.A. Gustafsson, K.S. Korach, D. Pfaff, S. Ogawa, Genotype/age interactions on aggressive behavior in gonadally intact estrogen receptor- α knockout (α ERKO) male mice, *Horm. Behav.* 41 (2002) 288–296.
- [87] S. Ogawa, E. Choleris, D. Pfaff, Genetic influences on aggressive behaviors and arousability in animals, *Ann. N. Y. Acad. Sci.* 1036 (2004) 257–266.
- [88] D. Pfaff, S. Ogawa, K. Kia, N. Vasudevan, C. Krebs, J. Frohlich, L.-M. Kow, Genetic mechanisms in neural and hormonal controls over female reproductive behaviors, in: D.W. Pfaff, A.P. Arnold, A.M. Etgen, S.E. Fahrbach, R.T. Rubin (Eds.), *Hormones, Behavior and Brain*, vol. 3, Academic Press, San Diego, 2002, pp. 441–510.